

BEHAVIORAL HEALTH IMPACTS OF COVID-19 **Workplace Trends, Resources, and Strategies**

Kira Mauseth, Ph.D.
Behavioral Health Strike Team

Agenda



**Defining
key terms**



**What to expect
from a behavioral
health standpoint
over the next few
months**



**Understanding
impacts to you and
your teams**




**Strategies for
increasing
resilience**

Definitions

- Burnout:** Exhaustion of body, mind, and motivation due to exposure to prolonged and unresolved work stress or frustration. Burnout is often a consequence of perceived disparity between the demands of the job and the resources that an employee has available to them.
- Compassion fatigue:** Emotional and physical exhaustion leading to a diminished ability to empathize or feel compassion for others, also described as secondary traumatic stress.
- Moral injury:** Strong feelings of guilt, shame, or anger due to not being able to provide the kind of care or service you want and expect to provide.
- Resilience:** The process – involving behaviors, thoughts, and actions – of adapting well in the face of adversity, trauma, tragedy, threats, or significant sources of stress. Can be developed by focusing on connection, purpose, and flexibility/adaptability.
- Resilience factors:** Conditions that help a person survive during and recover from a crisis or trauma- usually internal strengths and external resources.

Washington State Department of Health | 3




Key Things to Know

- **Three to four million** Washingtonians currently are experiencing *clinically significant* behavioral health symptoms.
 - Depression, anxiety, and acute stress are the most common.
 - Adolescents, teens, young adults, and older adults may need extra support.
- **Substance use related challenges are expected to continue.**
- **“Anniversary Reactions” will be varied and widespread**
- **Pandemic apathy will drive acting “out” and acting “in.”**
 - Acting “out”: Pretending like the pandemic no longer applies, refusing to comply with regulations, trying to act in a ‘business as usual’ capacity.
 - Acting “in”: Giving up on things getting back to normal, hopelessness and withdrawal, many symptoms consistent with major depressive disorder or depression generally.
- **An eventual return to baseline levels of functioning** for many people should occur around 14-18 months after the initial outbreak (May–July 2021), given the vaccine distribution timeline as an essential contributor to hope for many.


Washington State Department of Health | 5

Depression and Anxiety – WA adults




Washington State Department of Health | 6

Trauma Cascade Potential



Washington State Department of Health | 7

FIGURE 1. Weighted prevalence estimates* of self-reported stress and worry about psychosocial stressors among adults aged ≥18 years (N = 1,004), overall and by race/ethnicity† — Porter Novelli View 360 survey, United States, April and May 2020



Abbreviations: COVID-19 = coronavirus disease 2019; NH = non-Hispanic/Latino.
 * With 95% confidence intervals shown by error bars.
 † Other non-Hispanic minority groups include participants who identified as Native American/Alaska Native, Asian, multiracial, or another race/ethnicity.

FROM:
 Morbidity and Mortality Weekly Report 162
 MMWR / February 5, 2021 / Vol. 70 / No. 5 US
 Department of Health and Human
 Services/Centers for Disease Control and
 Prevention Racial and Ethnic Disparities in the
 Prevalence of Stress and Worry, Mental Health
 Conditions, and Increased Substance Use Among
 Adults During the COVID-19 Pandemic — United
 States, April and May 2020

Lela R. McKnight-Eily, PhD; Catherine A. Okoro,
 PhD; Tara W. Strine, PhD; Jorge Verlenden, PhD;
 NaTasha D. Hollis, PhD; Rashid Njai, PhD;
 Elizabeth W. Mitchell, PhD; Amy Board, DrPH;
 Richard Puddy, PhD; Craig Thomas, PhD

Washington State Department of Health | 8

Burnout, Compassion Fatigue, Moral Injury, and Exhaustion


Workplace burnout and similar phenomena continues to increase over time.

- Compounded by other factors, such as mental health stigma, PPE access, and added work.
- Burnout: Exhaustion of body and mind, unequal balance of demands and resources.
- Compassion fatigue: Emotional/physical tiredness, less ability to empathize.
- Moral injury: Strong feelings of guilt, shame, or anger due to not being able to provide the kind of care or service they want and expect to provide.


General fatigue, exhaustion, and feeling overwhelmed are common experiences.

- Sleep problems, diminished cognitive and high-level thinking, and increased impacts of existing behavioral health symptoms, such as depression, anxiety, or trauma.
- Organizations should address staff wellness and resilience, make it a priority, and model it.
- Practicing self-care, building personal coping/resilience plans, and rest are key for individuals.

Washington State Department of Health | 9



Washington State Department of Health | 10





The Good News

Typical long-term response to disasters is resilience, rather than disorder. Resilience is something that can be intentionally taught, practiced, and developed for people across all groups.

Resilience can be increased by:

- Focusing on developing social **connections**, big or small.
- Reorienting and developing a sense of **purpose**.
- Becoming **adaptive** and psychologically **flexible**.
- Focusing on **hope**.

Washington State Department of Health | 12



Communication Issues

- Remember how our pre-frontal cortex is influenced by stress.
- No one can have a logical *problem-solving* oriented talk when they are flooded.
- Take extra time to talk.
- Get space from difficult conversations, rather than continuing to push the issue.
- Check in on levels of rest/tiredness before pursuing important conversations.

Zones of Regulation

Color	Level of Alertness	Feelings
Blue	Low state of alertness	Bored, tired, sad, disappointed, sick, depressed, shy
Green	Perfect state of alertness	Happy, positive, thankful, proud, calm, content, ready to learn
Yellow	Higher state of alertness	Excited, silly, annoyed, worried, embarrassed, confused, nervous
Red	Too much alertness	Upset, angry, aggressive, mad, too excited, terrified, out of control



Washington State Department of Health | 15

Working with Anger and Hostility: **SAFE Model** for De-Escalation

Many people are expressing distress about the pandemic in an external way. This often manifests itself as anger. In order to de-escalate yourself and others, the **SAFE model** provides key concepts to keep in mind:

- S: Self
- A: Area Awareness
- F: Feelings
- E: Engagement

Washington State Department of Health | 16

SAFE: Self

- Tune in to yourself.
- Be aware of your own reactions, the tone of voice you use, your body language, and your choice of words.
- Monitor yourself in order to stay calm and to not take the situation personally, even if the words become personal.
- **Larger non-verbal messages are particularly important, particularly in the case where PPE (face masks) can interfere with people's ability to pick up on nuances of communication.**
- Be aware of the non-verbal things you are 'saying' to the other person. **Posture** and **Position** are important here.

Washington State Department of Health | 17

SAFE: Area Awareness

- Pay attention to your physical area.
- Notice the space and people around you.
- Your general physical area includes people, exits, (potential) weapons, available help, and other resources.
- **Don't position or keep yourself between an angry person and their exit.**

Washington State Department of Health | 18

SAFE: Feelings

- Employ active listening techniques to identify what the angry person is feeling underneath the anger.
- **Remember that anger is often related to other emotions like fear or sadness. It is easier for most people to direct emotions outwardly than deal with them internally.**
- By listening for feelings underneath anger, you can identify the cause of the emotions at the center of the issue.
- It is easier to empathize with someone who is angry when you understand what they may be afraid of or worried about.

Washington State Department of Health | 19

SAFE: Engagement

- If it is safe to do so, connect with the angry person by **engaging** to understand their story. Use active listening.
- Don't dismiss them or their concerns. Identify and **engage resources or other people or information** that may be able to address or help solve their problem or concern in some way.
- **Engage support for yourself** when you are in the position of dealing with an angry person or people.
- Don't keep a hostile interaction to yourself. Share it with others to get the support you need after dealing with a difficult person or situation.
- **Engage your resources (friends, family, social networks)** to increase your resilience.

Washington State Department of Health | 20

Resilience Development

Purpose

- What motivates you?
- What contributes to *compassion rewards*?
- What can you remind yourself of to help on a day-to-day basis (**don't think too long term or big picture**).

Connection

- How can you maintain existing connections with others?
- How can you develop new connections?
- **Connection can be anything that prevents isolation.**

Flexibility and Adaptability

- How can you be creative in physical distancing while leveraging connection?
- How can you adjust your physical space?
- How can you adapt your schedule to give yourself discreet and clear breaks and boundaries?

Hope

- What opportunities may exist where they didn't before?
- What are some surprise or hidden benefits that have come out of recent experiences?
- What examples do you have to shift your thinking from a 'threat' to a 'challenge'?

Washington State Department of Health | 21

Practice the **REST** Model

Reward: Reward yourself for a job well done. Build reinforcements into your work. Help pay attention to this aspect for maintaining resilience.

Establish: Establish healthy boundaries. When you are off duty, stick to that boundary.

Share: Share your feelings, concerns, and stories. Participate in support and consultation groups. Make time for connections and activities in your life.

Trust: Trust your support network and reach out as needed. Refer people elsewhere if you are too tired or compromised emotionally to be able to offer support.

Washington State Department of Health | 22

Resources

Training:

- Health Support Team (including train-the-trainer)
- PsySTART-Responder (frontline healthcare only)

Resources:

- MEDIC, REST, and SAFE models
- [Behavioral Health Group Impact Reference Guide](#)
 - Healthcare and behavioral health providers, outreach teams, post critical care individuals, etc.
 - Unique challenges and considerations
 - Support strategies (organizational, supervisory, and personal)
- Children and families: [Behavioral Health Toolbox for Families: Supporting Children and Teens During the COVID-19 Pandemic](#)
- Emergency and healthcare workers: [Coping During COVID-19 for Emergency and Healthcare Professionals](#)
- Businesses and workers: [COVID-19 Guidance for Building Resilience in the Workplace](#)

Washington State Department of Health | 23

Resources (cont.)

Webpages:

DOH – Forecasts, situation reports, guidance, and other resources:

- [Behavioral Health Resources Webpage](#)

State – General mental health resources and infographics:

- [Mental and Emotional Well-being Resources](#)
- [Infographic Library](#)

Looking for support?
Call Washington Listens at
1-833-681-0211



Washington State Department of Health | 24